

A New Short Chain Acetamide from the Biosphere and Bioactive Glycerolipids Extracted from the Marine Bivalve

Codakia orbicularis (Lucinidae)

Francesca Goudou *†¹, Olivier Grovel †², Ali Al-Mourabit ³,
Céline Moriou ³, Olivier Gros ¹ and Philippe Petit ¹

¹Institut de Systématique, Evolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles. Campus de Fouillole, 97110 Pointe-à-Pitre, France

²Faculty of Pharmacy, University of Nantes, EA 2160-Mer Molécules Santé, 9 rue Bias BP 61112, Nantes-cedex 1 44035, France

³Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Univ. Paris-Sud, Université Paris-Saclay, 1, av. de la Terrasse, 91198 Gif-sur-Yvette, France

(Received August 31, 2021; Revised October 28, 2021; Accepted October 29, 2021)

Abstract: The marine bivalve *Codakia orbicularis*, harbors endosymbiotic bacteria located in its gill filaments. Bioassay-guided fractionation of an ethyl acetate extract of the gill tissues led to antibacterial fractions (against Gram positive and negative bacteria). The fraction eluted in AcOEt/ MeOH 90:10 displayed the greatest bioactivity. This fraction was analyzed using usual chromatographic and spectrometric methods and revealed three compounds. The first one (compound **1**) was an acetamide isolated for the first time from the biosphere. The other two were mono-glycerolipids (compound **2**) and (compound **3**) with at the position sn-2 of their glycerol monounsaturated fatty acid chains (respectively C18:1 Δ^9 and C16:1 Δ^9). Compounds **2** and **3** showed bacteriostatic activity against two pathogenic Gram-negative bacteria using a disc diffusion assay method. These molecules have been isolated for the first time from a bivalve. Further investigations are currently in progress, so as to understand the role of these bioactive compounds into the regulation of *C. orbicularis*' endosymbionts and/or to free-living pathogens.

Keywords: Antibacterial activity; monoacylglycol; ceramide. © 2021 ACG Publications. All rights reserved.

1. Animal Source

Codakia orbicularis (Linné, 1758) individuals (40-60 mm shell length) were collected by hand from *Thalassia testudinum* sea-grass sediments in Guadeloupe (lat 16°09'01.7"N, long 61°33'42.9"W). These bivalves were identified by Pr. Olivier Gros (bivalve specialist).

* Corresponding author: E- Mail: francesca.goudou@univ-antilles.fr (F. GOUDOU).

† These authors have the same contribution to the work reported

2. Previous Studies

Codakia orbicularis is an edible tropical bivalve belonging to the Lucinidae family. In order to assess its edibility, the first chemical study on this lucinid (whole organism without shell) has been conducted by Berg and his co-workers [1]. In their work on the bivalve, no toxins were found confirming thus its edibility. Lipids were mostly reported as it is the case for most bivalves. Some years later, other studies mainly focused on understanding how such an organism can survive into a sulfidic habitat. Biological investigations proved that this lucinid harbors a single endosymbiotic sulfur-oxidizing bacterial species located in its gill filaments [2, 3]. Flow cytometric analysis has shown that these endosymbionts maintain the duplication of their genome within the bivalve's gills without cellular division (each bacterial cell harbors up to seven copies of their genomes) [4] suggesting the existence of bacteriostatic molecules able to control bacterial cell division [5]. From that time, a new lectine called codakine, has been identified in the gills of *C. orbicularis* by crystallography and microcalorimetry but was not bioactive [6]. Few years later, a new spiro-indolothiazine named orbicularisine [7] and acetylated nucleoside derivatives [8] have been characterized by our group using both chromatographic and spectroscopic methods but no antibacterial activity has been found for none of these molecules.

3. Present Study

In order to find potential bacteriostatic molecules from the endosymbiotic gills of *C. orbicularis*, a bio-guided chemical study has been led on the gills.

Bioactivity test-Agar diffusion test [9]: Bacterial strains were used for heterologous bioassay: *Escherichia coli* ATCC 35218, *Klebsiella pneumoniae* ATCC 700603, and *Pseudomonas aeruginosa* ATCC 27853, belonging to Gram negative bacteria; and *Enterococcus faecalis* ATCC 51299, *Micrococcus luteus* ATCC 10240, and *Streptococcus pneumoniae* ATCC 49619, belonging to Gram positive bacteria. Strains were cultivated on Mueller Hinton agar plate at 37°C for all bioassay experiments. Inocula were obtained using nutrient medium #1 (meat extract [5 g], agar [7.5 g], tryptic pepton [7.5 g], NaCl [2.5 g] and milliQ water [500 mL]; pH = 7) incubated at 37°C for 24 hours. One hundred microliters of a 10⁻² dilution culture were spread on an agar plate with nutrient agar medium #1. Concerning the antimicrobial assay on Gram negative bacteria, 50 µg-amoxicillin antibiotic disc (Biomérieux, France) was used as positive control while on Gram positive bacteria, 50 µg-vancomycin antibiotic disc was used. Sterile assay discs (5 mm-diameter, Dutscher) were placed by plate after spreading. One of them received twenty microliters of ethyl acetate (solvent used to dissolve the compounds) as negative control. For the remaining sterile discs, each received twenty microliters of the prepared solution. The plates were then incubated for 24 hours at 37°C. Inhibition of a target strain was determined as positive when the diameter of the inhibition zone was at least 2 mm greater than the negative control.

Bio-guided chemical study: Gill filaments of *C. orbicularis* were extracted with ethyl acetate, yielding twenty-five grams of crude extract. This crude extract proved to be active against two bacterial strains, *Escherichia coli* ATCC 35218 (Gram -) and *Micrococcus luteus* ATCC 10240 (Gram +), with inhibition halos of 15 mm and 10 mm respectively (see supplementary material, figure S19). The bioactive crude extract was chromatographed on silica gel column using a gradient of solvent mixtures starting from hexane/ethyl acetate followed by ethyl acetate/methanol to methanol. The fractions eluted with AcOEt/MeOH 90:10 V/V were the most active against the bacteria species mentioned earlier and were subjected to purification by HPLC. This latter purification led to the isolation of 0.5 mg of compound **1**, 2.5 mg of compound **2** and 2.3 mg of compound **3** (Figure 1). The structure of compounds **1-3** were elucidated by the current spectroscopic methods.

Compound 1: white solid; HRESIMS *m/z* 286.2390 [M+H]⁺ (calc. for 286.2377; C₁₆H₃₁NO₃); ¹H NMR (MeOD, 600 MHz) δ 5.69 (1H, td, *J* = 7.8/ 7.8/ 14.7 Hz, H-5), 5.47 (1H, dd, *J* = 7.3/ 14.7 Hz, H-4), 4.06 (1H, t_{large}, *J* = 7.3 Hz, H-3), 3.86 (1H, dd, *J* = 6.4/ 10.8 Hz, H-2), 3.67 (2H, m, H-1), 2.04 (2H, m, H-6), 1.95 (3H, s, H-16), 1.38 (2H, m, H-7), 1.29 (12H, m, H-8 to H-13), 0.90 (3H, t, *J* = 7.3 Hz, H-14); ¹³C NMR (MeOD, 150 MHz) δ 172.9 (C, C-15), 134.6 (CH, C-5), 131.0 (CH, C-4), 73.4 (CH, C-3), 62.2

A new short chain acetamide and bioactive glycerolipids

(CH₂, C-1), 56.9 (CH, C-2), 33.1 (CH₂, C-6), 30.3 (CH₂, C-7 to C-12), 23.7 (CH₂, C-13), 22.6 (CH₃, C-16), 14.3 (CH₃, C-14).

Compound 2: transparent oil; HRESIMS m/z 357.3009 [M+H]⁺ (calc. for 357.3005; C₂₁H₄₁O₄); ¹H NMR (CDCl₃, 600 MHz) δ 5.36 (2H, m, H-12 and H-13), 4.94 (1H, quin, J = 4.8 Hz, H-2), 3.85 (4H, d, J = 4.8 Hz, H-1 and H-3), 2.39 (2H, t, J = 7.5 Hz, H-5), 2.03 (4H, m, H-11 and H-14), 1.65 (2H, q, J = 7.1 Hz, H-6), 1.30 (20H, m, H-7 to H-10 and H-15 to H-20), 0.84 (3H, t, J = 7.3 Hz, H-21); ¹³C NMR (CDCl₃, 150 MHz) δ 174.4 (C, C-4), 130.0 (CH, C-12 and C-13), 74.9 (CH, C-2); 61.7 (CH₂, C-1 and C-3), 34.5 (CH₂, C-5), 27.3 (CH₂, C-11 and C-14), 25.0 (CH₂, C-6), 31.9 (CH₂, C-19), 29.5 (CH₂, C-7 to C-10 and C-15 to C-18), 22.7 (CH₂, C-20), 14.2 (CH₃, C-21).

Compound 3: transparent oil; HRESIMS m/z 329.2698 [M+H]⁺ (calc. for 329.2692, C₁₉H₃₇O₄); ¹H NMR (CDCl₃, 600 MHz) δ 5.33 (2H, m, H-12 and H-13), 4.89 (1H, quin, J = 4.8 Hz, H-2), 3.80 (4H, d, J = 4.8 Hz, H-1 and H-3), 2.35 (2H, t, J = 7.5 Hz, H-5), 2.00 (4H, m, H-11 and H-14), 1.62 (2H, q, J = 7.1 Hz, H-6), 1.29 (16H, m, H-7 to H-10 and H-15 to H-18), 0.87 (3H, t, J = 7.3 Hz, H-19); ¹³C NMR (CDCl₃, 150 MHz) δ 174.4 (C, C-4), 130.0 (CH, C-12 and C-13), 74.9 (CH, C-2), 61.7 (CH₂, C-1 and C-3), 34.5 (CH₂, C-5), 27.3 (CH₂, C-11 and C-14), 25.0 (CH₂, C-6), 31.9 (CH₂, C-17), 29.5 (CH₂, C-7 to C-10 and C-15 to C-16), 22.7 (CH₂, C-18), 14.2 (CH₃, C-19).




Figure 1. Structures isolated in the present work.

Compound **1** presented an exact mass of 285.2304 which suggested its molecular formula as C₁₆H₃₁NO₃ with two degrees of unsaturation (m/z 286.2390 [M+H]⁺; calc. 286.2377 $\delta \pm 0.0013$ ppm). Its ¹H NMR shown olefinic and aliphatic protons including a characteristic acetyl group observed by the H-16 proton at δ_H 1.95. From the ¹³C NMR data, it has been deduced that the only nitrogen into the structure of compound **1** was connected to the C-2 carbon at δ_C 56.9 which was bounded to a proton δ_H 3.86 (H-2). ¹H-¹H COSY spectra revealed that this H-2 proton was correlated to the H-3 methine proton at δ_H 4.06 and to the two H-1 methylene protons at δ_H 3.67. According to the ¹H-¹³C HSCQ spectra, the H-3 was supported by the C-3 carbon at δ_C 73.4 whereas the two H-1 by the C-1 carbon at δ_C 62.2. The chemical shifts of C-1 and C-3 carbons were characteristics of carbons linked to hydroxyl groups. The carbon C-3 at δ_C 73.4 from the 1,3-propanediol, 2-amino fragment found, have been connected via strong ¹H-¹³C HMBC correlations to the olefinic proton at δ_H 5.47 (H-4) corresponding to the carbon C-4 at δ_C 131.0 (¹H-¹³C HSQC data). The combined 2D NMR spectra revealed that the second carbon of the double bond, C-5 at δ_C 134.6 and wearing the H-5 proton (δ_H 5.69), was attached to a nine-carbon alkyl chain formed by aliphatic protons (H-6 – H-14) linearly assembled. Because of the large coupling between the olefinic protons (H-4; H-5, $J_{4,5}$ = 14.7 Hz) the *E* configuration was attributed to this double bond [10]. The above results led to a sphingoid base. Otherwise the ¹H-¹³C HMBC spectrum of compound **1** showed correlations between the protons at δ_H 1.95 ppm (H-16) and the C-15 carbonyl at δ_C 172.9 corresponding to an acetyl group (Figure 2). All these data shown that compound **1** was a *N*-acetylated sphingoid base. By comparing C-2 and C-3 chemical shifts with those of synthetic and natural ceramides, the relative configuration was deduced to be 2*S**, 3*R** which is a characteristic configuration in sphingosine-type bases of marine invertebrates [10, 11]. **1** is a 14-*ES* ceramide namely the *N*-[(2*S**, 3*R**, 4*E**)-3-hydroxy-2-(hydroxymethyl)-4-tridecen-1-yl]-acetamide. Natural ceramides usually occur with the configuration 2*S*, 3*R*, 4*E* [11]. Compound **1** is a natural ceramide formed by a C2 fatty acid and a C14 sphingosine backbone. Ceramides with short fatty acids (C2) have been only found before in mammals and are formed by an acetyl transfer from platelet-activating factor to a sphingosine [12]. The short chain acetamide (compound **1**) has been known until now only as a synthetic compound (CAS

number 2097561-20-9). Here we report the isolation of *N*-[(2*S**, 3*R**, 4*E**)-3-hydroxy-2-(hydroxymethyl)-4-tridecen-1-yl]-acetamide (compound **1**) for the first time from the biosphere.




Figure 2. ^1H - ^{13}C HMBC (arrows) and ^1H - ^1H COSY (bold) correlations in compound **1**.

The molecular formula of compound **2** was found to be $\text{C}_{21}\text{H}_{40}\text{O}_4$ ($\Delta m = 1,1$ ppm) by HRESIMS which displayed a protonated molecule $[\text{M} + \text{H}]^+$ at $m/z = 357.3009$. The HRESIMS of compound **3** gave a positive ion at $m/z = 329.2698$ corresponding to the molecular formula $\text{C}_{19}\text{H}_{36}\text{O}_4$ ($\Delta m = 1,8$ ppm). Compounds **2** and **3** shown a mass difference of 28 Da equivalent to two methylene groups and both molecules had similar ^1H NMR spectra showing a first part corresponding to signals of unsaturated fatty acids and a second part to a glycerol moiety [13, 14]. Monoglycerolipids with an oleic fatty acid or a palmitoleic fatty acid have been already discovered but the ^1H NMR spectra of compounds **2** and **3** showed some differences with those of the literature for the signals of the glycerol [15, 16, 17]. The ^1H NMR data of compound **2** are compared to those of an α -monoglyceride of oleic acid [16] in **Table 1**.

Table 1. ^1H NMR spectroscopic data of compound (**2**) compared to literature's data, both in CDCl_3 .

Present work (compound 2)			Literature (α -monoglyceride of oleic acid; [16])	
Location	Type	δ_{H} (J in Hz)	Type	δ_{H} (J in Hz)
12,13	CH	5.36, m	CH	5.35, m
2	CH	4.94, quin ($J = 4.8$ Hz)	CH	4.21, dd ($J = 11.7/ 4.3$ Hz)
1, 3	CH ₂	3.85, d ($J = 4.8$ Hz)	CH ₂	3.93, m; 4.15, dd ($J = 11.7/ 5.8$ Hz)
			CH ₂	3.60, dd ($J = 11.7/ 5.8$ Hz); 3.70, dd ($J = 11.7/ 3.9$ Hz)
5	CH ₂	2.39, t ($J = 7.5$ Hz)	CH ₂	2.35, t ($J = 7.6$ Hz)
11, 14	CH ₂	2.03, m	CH ₂	2.02, m
6	CH ₂	1.65, q ($J = 7.1$ Hz)	CH ₂	1.63, m
7-10 and 15-20	CH ₂	1.30, m	CH ₂	1.29, m
21	CH ₃	0.84, t ($J = 7.3$ Hz)	CH ₃	0.88, t ($J = 6.8$ Hz)
4	C	-----	C	-----

The analysis of both compounds was performed in deuterated chloroform. The signal differences between compound **2** and the α -monoglyceride of oleic acid from the literature have been presented in bold type in Table 1 and have been explained by the position of the fatty acid on the glycerol. In fact, the 4/1 ratio observed between the protons of the glycerol moiety at δ_{H} 4.94 (H-1 and H-3) and at δ_{H} 3.85 (H-2) proved an equivalence of these methylene and suggested a symmetry in the structure of compounds **2** and **3**. This symmetry was again validated by ^1H - ^{13}C HMBC and ^1H - ^{13}C HSQC correlations between the protons at δ_{H} 3.85 (H-2) and the carbon at δ_{C} 61.7 ($\text{CH}_{2\text{glycerol}}$). It was deduced that the fatty acid chain of compounds **2** and **3** was on the position *sn*-2 of the glycerol for both compounds' contrary to this one of the literatures which was on the position *sn*-1 (or *sn*-3). Furthermore, the multiplet at δ_{H} 5.36 (H-12 and H-13) indicated one unsaturation for each fatty acid and according to Liu and his co-workers [18] the chemical shifts of the associated carbons δ_{C} 27.3 (C-11 and C-14) confirmed an unsaturation with a *Z* configuration. In the literature, only D-9 mono-unsaturated fatty acids have been described as part of *sn*-2 substituted mono-glycerolipids isolated from plants or microalgae [19]. In addition to that, enzymatic hydrolysis of triglycerides often led to *sn*-2 mono-

A new short chain acetamide and bioactive glycerolipids extracted from a bivalve. As many kinds of glycerolipids are described to be bioactives [22], it encouraged glycerolipids [20] and the triglyceride 1-linoleoyl-2-oleoyl-3-stearoyl-glycerol has been previously isolated among 15 g of other triacylglycerols from *C. orbicularis* by our group (data not shown). We confronted our data to those given in the literature, which led us to the identification of compound **2** as being 9-octadecenoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester and compound **3** which was identified as 9-palmitoleic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester. So far, in the marine biosphere, monoglycerolipids have only been reported from algae [21] and sponges [15] but none have been previously the evaluation of the bioactivity of compounds **2** and **3** according to the agar diffusion method described above for assessing antibacterial activity.

The highest inhibition zone was obtained against *E. coli* ATCC 35218 with a diameter of 13.5 ± 0.5 mm for compound **2** and 12.5 ± 0.5 mm for compound **3**. Both compounds revealed a moderate activity against *K. pneumonia* ATCC 700603 (10.5 ± 0.5 mm). Compounds **2** and **3** were inactive against the Gram-positive bacteria tested but they showed an antibacterial activity against two Gram-negative bacteria. These results are interesting as 90% of the bacteria found into the marine biosphere are Gram-negative bacteria [23] and the positive bacteria detected in coastal marine environments are generally faecal contaminants of human origin or potentially human pathogens [24] with a short lifespan due to the salts. Glycerolipids have already presented several bioactivities (cyclooxygenase-II inhibitor; algicidal) [16]. Up to now, the dodecylglycerol was the only antibacterial natural product described in the literature [25]. Here, compounds **2** and **3** are reported as two new bioactive glycerolipids.

Acknowledgments

This work was supported by the Region Guadeloupe and the DEAL (Grant [11-026097-D]).

Supporting Information

Supporting Information accompanies this paper on <http://www.acgpubs.org/journal/records-of-natural-products>

ORCID

Francesca Goudou: [0000-0002-9979-6250](https://orcid.org/0000-0002-9979-6250)

Olivier Grovel: [0000-0003-3165-4771](https://orcid.org/0000-0003-3165-4771)

Ali Al-Mourabit: [0000-0002-1762-6910](https://orcid.org/0000-0002-1762-6910)

Céline Moriou: [0000-0003-2481-6461](https://orcid.org/0000-0003-2481-6461)

Olivier Gros: [0000-0002-2297-2009](https://orcid.org/0000-0002-2297-2009)

Philippe Petit: [0000-0002-5598-7557](https://orcid.org/0000-0002-5598-7557)

References

- [1] C. J. Berg, J. Krzynowek, P. Atalo and K. Wiggin (1985). Sterol and fatty acid composition of the clam, *Codakia orbicularis*, with chemoautotrophic symbionts, *Lipids* **20**, 116–120.
- [2] L. Frenkiel and M. Mouëza (1995). Gill ultrastructure and symbiotic bacteria in *Codakia orbicularis* (Bivalvia, Lucinidae), *Zoomorphology* **115**, 51–61.
- [3] O. Gros, A. Darrasse, P. Durand, L. Frenkiel and M. Mouëza (1996). Environmental transmission of a sulfur-oxidizing bacterial gill endosymbiont in the tropical lucinid bivalve *Codakia orbicularis*, *Appl. Environ. Microbiol.* **62**, 2324–2330.
- [4] A. Caro, O. Gros, P. Got, R. De Wit and M. Troussellier (2007). Characterization of the population of the sulfur-oxidizing symbiont of *Codakia orbicularis* (Bivalvia, Lucinidae) by single-cell analyses, *Appl. Environ. Microbiol.* **73**, 2101–2109.
- [5] A. Caro, P. Got, M. Bouvy, M. Troussellier and O. Gros (2009). Effects of long-term starvation on a host bivalve (*Codakia orbicularis*, Lucinidae) and its symbiont population, *Appl. Environ. Microbiol.* **75**, 3304–3313.
- [6] J. P. Gourdine and E. J. Smith-Ravin (2007). Analysis of a cDNA-derived sequence of a novel mannose-binding lectin, codakine, from the tropical clam *Codakia orbicularis*, *Fish Shellfish Immunol.* **22**, 498–509.

- [7] F. Goudou, P. Petit, C. Moriou, O. Gros and A. Al-Mourabit (2017). Orbicularisine: a new spiro-indolothiazine skeleton isolated from gills of the tropical bivalve *Codakia orbicularis*, *J. Nat. Prod.* **80**, 1693–1696.
- [8] F. Goudou, A. Al-Mourabit, C. Moriou, P. Dumesnil, O. Gros and P. Petit (2019). Acetylated nucleoside derivatives isolated from a shallow-water marine bivalve *Codakia orbicularis*, *Chem. Nat. Compd.* **55**, 790–792.
- [9] T. Brinkhoff, G. Bach, T. Heidorn, L. Liang, A. Schlingloff and M. Simon (2004). Antibiotic production by a *Roseobacter* clade-affiliated species from the German Wadden Sea and its antagonistic effects on indigenous isolates, *Appl. Environ. Microbiol.* **70**, 2560–2565.
- [10] V.P. Careaga and M.S. Maier (2014). Cerebrosides from marine organisms, *Stud. Nat. Prod. Chem.* **42**, 59–81.
- [11] J. S. O'Brien and G. Rouser (1964). The fatty acid composition of brain sphingolipids: sphingomyelin, ceramide, cerebroside, and cerebroside sulfate, *J. Lipid Res.* **5**, 339–342.
- [12] H. Van Overloop, Y. Denizot, M. Baes and P. P. Van Veldhoven (2007). On the presence of C2-ceramide in mammalian tissues: possible relationship to etherphospholipids and phosphorylation by ceramide kinase, *Biol. Chem.* **388**, 315–324.
- [13] W.W. Nawar (1996). Lipids, *Food Chem.* **1**, 225–319.
- [14] A. Cutignano, E. Luongo, G. Nuzzo, D. Pagano, E. Manzo, A. Sardo and A. Fontana (2016). Profiling of complex lipids in marine microalgae by UHPLC/Tandem mass spectrometry, *Algal Res.* **17**, 348–358.
- [15] Y. Liu, J. H. Jung, H. Ji and S. Zhang (2006). Glycerolipids from a *Sarcotragus* species sponge, *Molecules* **11**, 714–719.
- [16] S. Hirao, K. Tara, K. Kuwano, J. Tanaka and F. Ishibashi (2012). Algicidal activity of glycerolipids from brown alga *Ishige sinicola* toward red tide microalgae, *Biosci. Biotechnol. Biochem.* **76**, 372–374.
- [17] Y. Sun, H. Wang, G. Guo, Y. Pu, B. Yan and C. Wang (2015). Isolation, purification, and identification of antialgal substances in green alga *Ulva prolifera* for antialgal activity against the common harmful red tide microalgae, *Environ. Sci. Pollut. Res.* **23**, 1449–1459.
- [18] Y. Liu, C. Lee, J. Hong and J. H. Jung (2002). Cyclitol derivatives from the sponge *Sarcotragus* species, *Bull. Korean Chem. Soc.* **23**, 1467–1469.
- [19] J. S. Cheng, Y. H. Niu, S. H. Lu and Y. J. Yuan (2012). Metabolome analysis reveals ethanolamine as potential marker for improving lipid accumulation of model photosynthetic organisms, *J. Chem. Technol. Biotechnol.* **87**, 1409–1418.
- [20] L. Y. Yang and A. Kuksis (1991). Apparent convergence (at 2-Monoacylglycerol level) of phosphatidic acid and 2-monoacylglycerol pathways of synthesis of chylomicron triacylglycerols, *J. Lipid Res.* **32**, 1173–1186.
- [21] H. W. Chang, K. H. Jang, D. Lee, H. R. Kang, T. Y. Kim, B. H. Lee, B. W. Choi, S. Kim and J. Shin (2008). Monoglycerides from the brown alga *Sargassum sagamianum*: isolation, synthesis, and biological activity, *Bioorg. Med. Chem. Lett.* **18**, 3589–3592.
- [22] E. D. Costa, J. Silva, S. Mendonça, M. Abreu and M. Domingues (2016). Lipidomic approaches towards deciphering glycolipids from microalgae as a reservoir of bioactive lipids, *Mar. Drugs* **14**, 101.
- [23] W. Fenical (1993). Chemical studies of marine bacteria: developing a new resource, *Chem. Rev.* **93**, 1673–1683.
- [24] M. M. Lleò, C. Signoretto and P. Canepari (2005). Gram-Positive bacteria in the marine environment, In: *Oceans and Health: Pathogens in the marine environment*, eds: S. Belkin, R.R. Colwell, Springer, Boston, Massachusetts, pp.307–330.
- [25] H. S. Ved, E. Gustow, V. Mahadevans and R. A. Pieringer (1984). Dodecylglycerol: a new type of antibacterial agent which stimulates autolysin activity in *Streptococcus faecium* ATCC 9790, *J. Biol. Chem.* **259**, 8115–8121.